

Discussion of track structure models and efficiencies of luminescence dosimeters to HCP

Gabriel O. Sawakuchi and Eduardo G. Yukihiro

*Oklahoma State University, Department of Physics
Technology & Research Park – Venture I, 1110 S. Innovation Way
Stillwater, OK 74075, USA*

In this work we present a discussion on preliminary predictions of efficiencies to HCP using four different analytical track structure models: Chatterjee [1,2], Katz [3,4], Katz-Waligorski [5] and Katz-Fageeha [6]. The method used in this work consists of using an analytical track structure model convoluted with the dose response of the luminescence material.

1.1 Chatterjee's model

In Chatterjee's model (equation 1.1) [1,2] $D(r)$ is the local dose at a radial distance r from the path of the primary particle, r_c is the radius of the so called core region and r_p is the radius of the penumbra region. Expressions for r_c and r_p are given together with equation 1.1.

1. Track Structure

1.1 Chatterjee's model

$$D(r) = \left[\begin{array}{l} \frac{LET_{\infty} / 2}{r_c^2} + \frac{LET_{\infty} / 2}{r_c^2 \left(1 + 2 \ln \frac{r_p}{r_c} \right)}, \quad r < r_c = \sqrt{\frac{4 n e^2}{m}} v \\ \frac{LET_{\infty} / 2}{r_c^2 \left(1 + 2 \ln \frac{r_p}{r_c} \right)}, \quad r_c < r < r_p = 396 v^{2.7} \end{array} \right] \quad (1.1)$$

1.2 Katz's model

$$D(r) = \frac{Ne^4}{mc^2} \frac{Z^{*2}}{\alpha \beta^2} \frac{1}{r} \left[\frac{1 - \left(1 - \frac{r + \theta}{R + \theta}\right)^{1/\alpha}}{r + \theta} \right] \quad (1.2)$$

where

$$R = kW^\alpha; \quad \theta = kI^\alpha$$

k and α are constants, I is the ionization potential, W is the maximum delta ray energy [3,4]

$$W = 2mc^2 \frac{\beta}{1 - \beta^2}$$

1.3 Katz-Waligorski's model

In order to correct Katz's model to fit to the experimental data in the region of small radial distances and also to predict the correct stopping power when the distribution of dose is radially integrated, Waligorski *et al.* [5] multiplied equation 1.2 by an arbitrary function.

$$D(r) = D_{Katz}(r) [1 + K_w(r)]$$

1.3 Katz-Waligorski's model

$$K_w(r) = A \left(\frac{r-B}{C} \right) \exp \left(-\frac{r-B}{C} \right) \quad (1.3)$$

where

$$(a) \quad r > B = 0.1 \text{ nm}, \quad C = 1.5 \text{ nm} + 5 \text{ nm} \times \beta$$

$$A = \begin{bmatrix} 8 \times \beta^{1/3}, & \text{for } \beta < 0.03 \\ 19 \times \beta^{1/3}, & \text{for } \beta > 0.03 \end{bmatrix}$$

$$(b) \quad r < B = 0.1 \text{ nm}$$

$$K_w(r) = 0$$

1.3 Katz-Fageeha's model

Fageeha *et al.* [6] improved Katz-Waligorski's model using a different arbitrary function

$$K_F(r) = A \beta^B (r - 0.1) \exp\left(-\frac{r}{C}\right)$$

where

$$B = 0.215; \quad C = 3.127 - 0.434 \beta$$

$$A = \begin{bmatrix} 0, & \text{for } \beta < 0.0081 \\ 112 \beta - 0.899, & \text{for } 0.008 < \beta < 0.09 \\ 0.674 \beta + 9.21, & \text{for } \beta > 0.091 \end{bmatrix}$$

2.2 Experimental data




Figure 1: Dose response of luxel irradiated with a ^{90}Sr source.

3.2 Results

$$\eta = \frac{\int_{r_{min}}^{r_{max}} r S[D(r)] dr}{\int_{r_{min}}^{r_{max}} r S_0[D(r)] dr}$$

Table 1: Comparison between experimental efficiencies of luxel and predictions using different track structure models.

Beam	LET (keV/ m)	Experiment	Chatterjee	Katz	Katz-Waligorski	Katz-Fageeha
H: 230 MeV	0.41	0.951(6)	0.80	0.57	0.81	0.83
H: 70 MeV	0.96	0.928(2)	0.53	0.92	0.92	0.91
He: 150 MeV/n	2.18	0.923(7)	0.53	0.86	0.88	0.84
C: 400 MeV/n	10.96	0.684(7)	0.50	0.74	0.76	0.64
C: 290 MeV/n	12.97	0.626(8)	0.49	0.79	0.76	0.63
O: 400 MeV/n	19.48	0.609(7)	0.48	0.75	0.72	0.57
Ne: 400 MeV/n	30.44	0.467(16)	0.47	0.75	0.68	0.52
Fe: 500 MeV/n	186.3	0.341(20)	0.43	0.71	0.50	0.35

- All the models failed to predict efficiencies of high LET_∞ particles where the efficiency levels off (Table 1 and Figure 2).
 - Chatterjee and Katz models are very poor to predict the efficiencies (Table 1).
 - Katz-Waligorski and Katz-Fageeha models fail to predict efficiencies of high energy particles. The models give unphysical results for energies higher than a limit that depend on the type of particle (see proton data).
-

- Although Katz-Waligorski and Katz-Fageeha models have deficiencies, the models presented reasonable results for the type of particles and energies studied here (Figure 2).




Figure 2: Comparison between experimental efficiencies and predictions using Katz-Waligorski and Katz-Fageeha models.

OSU 4. Other points to consider

- A “monoenergetic” dose response curve may not reproduce the delta ray energy spectrum created when a HCP transverses the material.
 - The fitted range-energy relation used by Katz, Katz-Waligorski and Katz-Fageeha models deviates from experimental data for electrons of energies larger than 1 MeV [5].
-

- [1] A. Chatterjee and H. J. Schaefer, “Microdosimetric structure of heavy ion tracks in tissue”, *Radiat. Environ. Biophys.* **13**, 215-227 (1976).
- [2] A. Chatterjee and W. R. Holley, “Computer simulation of initial events in the biochemical mechanisms of DNA damage”, *Adv. Radiat. Biol.* **17**, 181-226 (1993).
- [3] J. J. Butts and R. Katz, “Theory of RBE for heavy ion bombardment of dry enzymes and viruses”, *Radiat. Res.* **30**, 855-871 (1967).
- [4] Zhang Chunxiang, D. E. Dunn and R. Katz, “Radial distribution of dose and cross-sections for the inactivation of dry enzymes and viruses”, *Radiat. Prot. Dosim.* **13**, 215-218 (1985).
- [5] M. P. R. Waligórski, R. N. Hamm and R. Katz, “The radial distribution of dose around the path of a heavy ion in liquid water”, *Nucl. Tracks Radiat. Meas.* **11**, 309-319 (1986).
- [6] O. Fageeha, J. Howard and R. C. Block, “Distribution of radial energy deposition around the track of energetic charged particles in silicon”, *J. Appl. Phys.* **75**, 2317-2321 (1993).
-

The experiments were performed as part of the ICCHIBAN international inter-comparison project using heavy ions at NIRS-HIMAC. The authors thank Dr. Y. Uchihori, Dr. N. Yasuda, Hisashi Kitamura and Dr. E. Benton for their support in irradiations at NIRS-HIMAC.
